CRANIUM-Interactions with CPHS Formulary Psychiatric Drugs* January 14, 2015

No absolute contraindications with any of these medications

METFORMIN

Antidepressants

- Bupropion (Wellbutrin): Metformin may increase serum concentration of buproprion
- Fluoxetine (Prozac): May increase risk of hypoglycemia with metformin

Mood Stabilizers

Lamotrigine (Lamictal): Metformin may increase serum concentration of lamotrigine

Other

- Atenolol (Tenormin) and Propranolol (Inderal): May alter glucose metabolism, prolong hypoglycemia, mask hypoglycemic symptoms
- Levothyroxine (T4, Synthroid) & Liothyronine (T3, Cytomel): May decrease hypoglycemic efficacy of metformin
- Nicotine replacement: Nicotine may increase metformin levels and risk of lactic acidosis

AMLODIPINE (Norvasc)

Antipsychotics

Aripiprazole (Abilify): Amlodipine may increase the serum concentration of aripiprazole

Antidepressants

- Duloxetine (Cymbalta) & Flucoxamine (Luvox): Amlodipine may increase orthostatic hypotensive effect of these medications
- MAO Inhibitors: May increase the hypotensive efficacy of amlodipine.

Mood Stabilizers

 Carbamazepine (Tegretol) & oxcarbazepine (Trileptal): May decrease the hypotensive efficacy of amlodipine

Other

• Methylphenidate (Ritalin): May diminish the antihypertensive efficacy of amlodipine

ATORVOSTATIN (Lipitor)

Antipsychotics

Aripiprazole (Abilify): Atorvostatin may increase the serum concentration of Aripiprazole

Antidepressants

Fluoxetine (Prozac), Fluvoxamine (Luvox), and Nefazodone (Serzone): May increase statin level and increase risk of myopathy and rhabdomyolysis

Mood Stabilizers

• Carbamazepine (Tegretal) & Oxcarbazepine (Trileptal): May diminish antihyperlipidemic efficacy

Other

- Atenolol (Tenormin) and Propranolol (Inderal): May decrease antihyperlipidemic efficacy
- Naloxegol (Movantic): Atorvostatin may increase concentration of naloxegol.